


Codeine content in Medications

- 8 mg Codeine in each solpadeine tablet
- 12.8mg Codeine in each Nurofen Plus tablet
- 15mg Codeine in each in Codipar tablet
- 30mg Codeine in each Tylex/Kapake/Solpadol tablet
- Advice to patients:
 - We know codeine is not a good analgesic
 - We know it causes constipation and is habit forming





One Difference between Codeine and Morphine





Just a methyl group!



NSAIDs



- Responsible for 30% of ADRs
 - Bleeding, MI, stroke and nephrotoxicity
- GI bleeding risk increased from day 1 (and more likely fatal vs GI bleed from H. Pylori)
 - Increased 12 fold if co-prescribed with corticosteroids
 - Increased 11 fold if co-prescribed with spironolactone
 - SSRIs 7 fold



BJGP 2016;66;172





Adverse effects of NSAIDs

- · Increased risk of MI and Stroke
 - Reducing Cox-2
- Reduces antiplatelet effect of aspirin
- Nephrotoxicity
 - Do renal function test on patients who take ibuprofen regularly
 - Doubles rate of AKI if over 65 years



BJGP 2016;66;172





Adverse effects of NSAIDs

- · Increased risk of MI and Stroke
 - Reducing Cox-2
- Reduces antiplatelet effect of aspirin
- Nephrotoxicity
 - Do renal function test on patients who take ibuprofen regularly
 - Doubles rate of AKI if over 65 years

BJGP 2016;66;172

