




# Nuvaxovid® (Novavax) Summary Sheet

please refer to [www.immunisation.ie](http://www.immunisation.ie) for full guidance

**This vaccine may be given to people aged 18 years and older who have a contraindication to an mRNA vaccine, or who have declined another COVID-19 vaccine**



## Storage

- +2°C to +8°C until expiry date
- Must be used within 6 hours from first puncture ("discard" date and time)

## Primary course

- 0.5ml dose
- 2 doses 21 days apart
- People aged 18 years and older

## Additional dose\*

- 0.5ml dose
- 2 months after primary course
- People aged 18 years and older and immunocompromised

## Booster dose\*

- 0.5ml dose
- 5 months after the last dose
- People aged 18 years and older

## Preparation

### Inspect



- Remove the vaccine from fridge
- Check "expiry" date
- Inspect vial
- Gently swirl the vial

### Label the vial



- Label the vial with "discard time and date" (6 hours after time of first puncture)

## Administration



Check the "discard" date and time




Clean top of vial with alcohol swab



Withdraw dose 0.5ml of vaccine



Administer dose intramuscularly



Version 1.0 22 February 2022

Created by: HSE National Immunisation Office

\*Do not give if the person received 2 doses of Nuvaxovid® as the primary course (a homologous Nuvaxovid® primary course)